CLAY COUNTY REPORT OF ENDANGERED, THREATENED, AND SPECIAL CONCERN PLANTS, ANIMALS, AND NATURAL COMMUNITIES OF KENTUCKY

PRESERVES COMMISSION 801 SCHENKEL LANE FRANKFORT, KY 40601 (502) 573-2886 (phone) (502) 573-2355 (fax)

www.naturepreserves.ky.gov

Kentucky State Nature Preserves Commission Key for County List Report

Within a county, elements are arranged first by taxonomic complexity (plants first, natural communities last), and second by scientific name. A key to status, ranks, and count data fields follows.

STATUS

KSNPC: Kentucky State Nature Preserves Commission status:

USESA: U.S. Fish and Wildlife Service status:

SOMC = Species of Management Concern

RANKS

GRANK: Estimate of element abundance on a global scale:

G1 = Critically imperiled GU = Unrankable

G2 = Imperiled G#? = Inexact rank (e.g. G2?)
G3 = Vulnerable G#Q = Questionable taxonomy

G4 = Apparently secure G#T# = Infraspecific taxa (Subspecies and variety abundances are coded with a 'T' suffix; the 'G'

G5 = Secure portion of the rank then refers to the entire species)

GH = Historic, possibly extinct GNR = Unranked GX = Presumed extinct GNA = Not applicable

SRANK: Estimate of element abundance in Kentucky:

S1 = Critically imperiled SU = Unrankable Migratory species may have separate ranks for different

S2 = Imperiled S#? = Inexact rank (e.g. G2?) population segments (e.g. S1B, S2N, S4M):

S3 = Vulnerable S#Q = Questionable taxonomy S#B = Rank of breeding population
S4 = Apparently secure S#T# = Infraspecific taxa S#N = Rank of non-breeding population
S5 = Secure SNR = Unranked S#M = Rank of transient population

SH = Historic, possibly extirpated SNA = Not applicable

SX = Presumed extirpated

COUNT DATA FIELDS

OF OCCURRENCES: Number of occurrences of a particular element from a county. Column headings are as follows:

- E currently reported from the county
- H reported from the county but not seen for at least 20 years
- F reported from county & cannot be relocated but for which further inventory is needed
- X known to be extirpated from the county
- U reported from a county but cannot be mapped to a quadrangle or exact location.

The data from which the county report is generated is continually updated. The date on which the report was created is in the report footer. Contact KSNPC for a current copy of the report.

Please note that the quantity and quality of data collected by the Kentucky Natural Heritage Program are dependent on the research and observations of many individuals and organizations. In most cases, this information is not the result of comprehensive or site-specific field surveys; many natural areas in Kentucky have never been thoroughly surveyed, and new species of plants and animals are still being discovered. For these reasons, the Kentucky Natural Heritage Program cannot provide a definitive statement on the presence, absence, or condition of biological elements in any part of Kentucky. Heritage reports summarize the existing information known to the Kentucky Natural Heritage Program at the time of the request regarding the biological elements or locations in question. They should never be regarded as final statements on the elements or areas being considered, nor should they be substituted for on-site surveys required for environmental assessments.

KSNPC appreciates the submission of any endangered species data for Kentucky from field observations. For information on data reporting or other data services provided by KSNPC, please contact the Data Manager at:

Kentucky State Nature Preserves Commission 801 Schenkel Lane Frankfort, KY 40601 phone: (502) 573-2886 fax: (502) 573-2355

email: naturepreserves@ky.gov internet: www.naturepreserves.ky.gov

County	/ Taxonomic Group	Scientific name	Common name	Statuses	Ranks		# of	Occi	urrer	ices
	Habitat					E	Н	F	Χ	U
Clay	Vascular Plants Moist or dry sandy woods, meador	Hypericum crux-andreae ws and barrens. also pine flatwoods (Weakley 1998).	St. Peter's-wort	Т/	G5 / S2S3	0	1	0	0	0
Clay	Vascular Plants MESIC WOODED RAVINES AND	Juglans cinerea ALONG STREAMS	White Walnut	S/SOMC	G3G4 / S3	1	0	0	0	0
Clay	Vascular Plants DRY TO MESIC SLOPE AND BO	Lathyrus venosus TTOMLAND FORESTS AND WOODLANDS, ESPECIALLY	Smooth Veiny Peavine IN BASE-RICH SOILS (WEAKLEY 1998).	S/	G5 / S2S3	1	0	0	0	0
Clay	Vascular Plants SWIFTLY FLOWING WATER, AT	Podostemum ceratophyllum TACHED TO ROCKS IN RAPIDS OF LARGER RIVERS	Threadfoot	S/	G5 / S3	12	0	0	0	0
Clay	Vascular Plants Calcareous forests and thickets us	Prenanthes crepidinea sually in alluvial areas.	Nodding Rattlesnake-root	Τ/	G4 / S2	4	0	0	0	0
Clay	Vascular Plants DRY- MESIC (CLOSED OR OPEN	Silphium wasiotense N) WOODLANDS AND ADJ. ROADSIDES AND A RAVINE I	Appalachian Rosinweed N MIX MESOPHYTIC FOREST.	S/SOMC	G3? / S3?	27	0	2	0	0
Clay	Vascular Plants Rich or open woods, chiefly in the	Solidago curtisii uplands; base of bluffs and along bluff ledges (Steyermark 1	Curtis' Goldenrod 1975).	Т/	G4G5 / S2S3	1	0	0	0	0
Clay	Vascular Plants Dry wood slopes and ridges.	Thermopsis mollis	Soft-haired Thermopsis	E/	G3G4 / S1	4	0	0	0	0
Clay		Anguispira rugoderma THE NORTH SIDE OF PINE MOUNTAIN (HUBRICHT 1985) COOL, BUT BURROWS INTO ROTTING WOOD AND SOIL			G2 / S2 PRING AND FALL WH	1 EN	0	0	0	0
Clay	Gastropods UNDER LEAF LITTER ON WOOD	Mesomphix rugeli DED HILLSIDES OR ON MOUNTAINS (HUBRICHT 1985).	Wrinkled Button	Τ/	G4 / S2	3	0	0	0	0
Clay	Freshwater Mussels INHABITS SAND, SILT, MUD, AN STREAMS.	Anodontoides denigratus D SMALL GRAVEL OFTEN NEAR COBBLE AND BOULDE	Cumberland Papershell RS IN POOLS AND RUNS WITH SLOW CUI	E / SOMC RRENT IN SMALL	G1 / S1 TO MEDIUM-SIZED	1	0	0	0	0
Clay	Freshwater Mussels	Epioblasma triquetra	Snuffbox	E / SOMC	G3 / S1	6	2	1	0	0
		o large rivers generally on mud, rocky, gravel, or sand subst oly buried in substrate and overlooked by collectors.	rates in flowing water (Baker 1928, Buchanar	n 1980, Johnson 1	978, Murrary and Leon	ard				
Clay	Freshwater Mussels GRAVEL BARS AND DEEP POOL ALLEN 1964, PARMALEE 1967).	Fusconaia subrotunda subrotunda LS IN LARGE RIVERS AND LARGE TO MEDIUM-SIZED S	Longsolid TREAMS (AHLSTEDT 1984, GOODRICH AN	S / ID VAN DER SCH.	G3T3 / S3 ALIE 1944, NEEL AND	2	0	0	0	0
Clay	Freshwater Mussels INHABITS SMALL TO MEDIUM-S	Villosa lienosa IZED RIVERS, USUALLY IN SHALLOW WATER ON A SAN	Little Spectaclecase ND/MUD/DETRITUS BOTTOM (PARMALEE	S / 1967, GORDON A	G5 / S3S4 ND LAYZER 1989).	5	2	0	0	0
Clay	Crustaceans ROCKY STREAMS (HOBBS 1989	Cambarus parvoculus	Mountain Midget Crayfish	Τ/	G4 / S2	3	0	0	0	0
Clay	• • •	Dryobius sexnotatus ix hardwood forest habitat, where it principally lives on sugar e typically found (Mike Bratton, pers comm).	Sixbanded Longhorn Beetle maple and, to a lesser extent, beech and elr	T / SOMC n (Perry et al. 1974	GNR / S1 4, Schweitzer 1989). M	1 id	0	0	0	0
Clay	Insects SAND AND GRAVEL IN SWIFTLY	Ophiogomphus howei Y FLOWING, UNPOLLUTED AND UNDAMMED RIVERS (C.	Pygmy Snaketail ARLE 1987, COOK 1992).	T/SOMC	G3 / S1S2	0	1	0	0	0

Data Current as of February 2006

County Report of Endangered, Threatened, and Special Concern Plants, Animals, and Natural Communities of Kentucky Kentucky State Nature Preserves Commission

County	/ Taxonomic Group	Scientific name	Common name	Statuses	Ranks		# of	Осс	urrer	ices
	Habitat					Е	Н	F	X	U
Clay		Ichthyomyzon fossor ND STREAMS WHERE ADULTS LIVE IN IS REQUIRE MIXED SAND, SILT, AND D	Northern Brook Lamprey SAND-GRAVEL BOTTOMS OF CLEAN RIFFLES AND RACE EBRIS IN QUIET WATER.	T / EWAYS (BURR ANI	G4 / S2 D WARREN 1986, PAC	0 SE	1	0	0	0
Clay	clearcuts, highway and powerline	rights-of-way (Hulse et al. 2001), rocky blu	Coal Skink ter and loose rocks; often the lizard occurs in the vicinity of spr uffs above creek valleys, dry, rocky, south-facing hillsides (Joh nes they take refuge in water. One nest was under a piece of s	nson 2000), and dr	y shale barrens (West	1 ts	0	0	0	0
Clay		· · · · · · · · · · · · · · · · · · ·	Sharp-shinned Hawk DUS, PRIMARILY IN CONIF. IN MORE NORTHERN AND MO IG RIDGES, LAKESHORES, & COASTLINES (B83NAT01NA)		G5 / S3B,S4N FION OF RANGE (B83	1	0	0	0	0
Clay	Breeding Birds Grasslands and savanna, especia	Cistothorus platensis Ily where wet or boggy, sedge marshes, k	Sedge Wren ocally in dry cultivated grainfields. In migration and winter also	S / in brushy grassland	G5 / S3B ls. (B83COM01NA)	0	1	0	0	0
Clay	Mammals Rafinesque's big-eared bats use a buildings, etc. Apparently less fred	,	Rafinesque's Big-eared Bat es, protected sites along clifflines, old mine portals, abandoned	S / SOMC I tunnels, cisterns, o	G3G4 / S3 old or seldom used	3	0	0	0	0

Data Current as of February 2006 Page 5 of 5